5.30 a.m. (Darwin time) when the ship was off Pt. Blaze, Northern Territory, and 58 miles offshore, a Nankeen Kestrel (Falco cenchroides) landed in an exhausted eondition. It fluttered about the ship and after being eaught was placed in a box with water. Next morning after being fed it was released. After eireling the ship it alighted on the mast. It was not seen on May 16 or 17 but reappeared again on the 18th, when it was seen sitting on the mast. It eontinually showed up throughout the trip of 15 days which took us to the southern end of the 80 Mile Beach and Onslow and back to Darwin. The Kestrel is known to migrate between Australia and Indonesia.

-ROBERT BOYD, Daglish.

Fan-tailed Cuckoo parasitising Western Silvereye.—On November 11, 1952, at Coolbellup Lake, 6 miles east of Fremantle, I found a nest of the Western Silvereye (Zosterops gouldi) eontaining an egg of the Fan-tailed Cuekoo (Cacomantis pyrrhophanus), with two of the host's eggs. The euckoo's egg measured 21.3 x 15.2 mm. and was dull white, marked with minute brownish spots, with a few underlying markings of light grey, which tended to form a zone around the broader end. The nest was in a small bush (Regelia ciliata), about 5 ft. from the ground.

This euckoo is a very searee species in this neighbourhood and this is the only occasion on which I have found its egg locally. It is also the only occasion on which I have found the Silvereye parasitised by any euckoo.

The Fan-tailed Cuekoo almost invariably chooses as hosts birds which build domed nests, but there are rare records in the Eastern States of its eggs being found in eup-shaped nests (including honeyeaters, robins and sittellas). The example I report appears to be the first in which the Western Silvereye has been parasitised.

-JULIAN R. FORD, Fremantle.

Dispersal Opportunities for Exotic Snails.—Two instances recently eams under notice at the Fremantle waterfront on the means by which the European snails, *Theba pisana* and *Helix aspersa*, might extend their range in Western Australia. Early in October, 1956, M.V. *Taranaki* discharged steel and general eargo from United Kingdom ports, Adhering to a girder was found an adult, 4-whorled specimen of *Helix aspersa* which, when moistened, quickly became active. Such snails can easily become secreted in eargo and generally distributed.

About three years ago the State ship *Dulverton* lifted deek eargo of jetty piles for a North-West port. The butts of most piles were deeply split, providing refuge for a large number of aestivating *Theba pisana*. The piles had lain for some time in the old Public Works Department yard, now the site of No. 10 berth of the North Wharf, which was heavily overrun with snails. Without doubt reports of the species' appearance near our North-West ports may soon be expected.

—G. KENDRICK, Fremantle.